

Medication Use Among Those With Osteoporosis in the Community

Tiffany Gill¹, Patrick Phillips², Debra Rowett³, Anne Taylor¹, Narsing Laddipeerla², and the North West Adelaide Health Study Team

¹ Population Research and Outcome Studies Unit, SA Health, Adelaide, South Australia, Australia

² Endocrinology Unit, The Queen Elizabeth Hospital, Woodville, South Australia, Australia

³ Drug and Therapeutic Information Service, Repatriation General Hospital, South Australia, Australia

INTRODUCTION

As the population ages the prevalence of osteoporosis and fractures increases. Medications form part of the treatment for osteoporosis, however, some may be missing (that is medications not taken), while others may be inappropriate. The aim of this study was to determine the prevalence of osteoporosis and medication use in a population-based cohort study conducted in Australia.

METHODS

The North West Adelaide Health Study is a representative longitudinal cohort study of people aged 18 years and over living in the north west region of Adelaide. The original sample (n=4060) was randomly selected and recruited by telephone interview to participate in a clinic assessment. In Stage 2 of the study (2004-06), information regarding the prevalence of musculoskeletal conditions was included.

Osteoporosis and Medications

The self-reported prevalence of osteoporosis and falls were collected using a Computer Assisted Telephone Interview (CATI) survey and fracture information from the self-completed questionnaire. All medications that participants were taking, including complementary and alternative medicines, were recorded at the clinic visit. Those aged 50 years and over who attended the clinic were offered the opportunity to have a DXA scan.

Stage 1

In Stage 1, all households with a telephone connected and the telephone number listed in the Electronic White Pages were eligible for random selection. Within each household, the person who had their birthday last and was aged 18 years or older, was selected for interview and invited to attend the study clinic. Of those interviewed (n=5850), 69% participated in the clinic visit (n=4060). Data were obtained from self-reported questionnaires and biomedical measurements.

Stage 2

Of the 4060 participants in Stage 1, 100 had died, leaving 3960 to be contacted, of which 81% (n=3206) attended the Stage 2 clinic (Figure 1).




Figure 1. Number of participants in the North West Adelaide Health Study

RESULTS

There were n=3502 participants who responded to the CATI component of the study. Overall, 3.8% (95% CI 3.2-4.5) and 8.8% (95% CI 7.5-10.4) of those aged 50 years and over, reported that they had been told by a doctor that they had osteoporosis.

Among respondents who had a fracture as a result of a fall from a standing height or less in the last five years, 16.8% (95% CI 12.2-22.6) had been told by a doctor that they had osteoporosis and 12.0% were on benzodiazepines, medications that are known to increase the risk of falls.

Of the participants who underwent a DXA scan, 3.6% (95% CI 2.6-4.9) had osteoporosis as defined by a T-score of -2.5 or less and 15.0% (95% CI 13.0-17.3) were osteopenic (T-score less than -1.0 and greater than -2.5). Of the participants 50 years and over with self-reported osteoporosis who had a DXA scan only 9.0% had a bone mineral density in the osteoporotic range.

Of those reporting that they had been told they have osteoporosis, 43.9% used bisphosphonates, with alendronate the most commonly used (65.4%, Figure 2).




Figure 2. Proportion taking each type of bisphosphonate

However, only 12.1% of those stating they had osteoporosis and taking a bisphosphonate also took both calcium and Vitamin D. Of those with osteoporosis and taking a bisphosphonate, 31.4% were also taking proton pump inhibitors which might suggest gastroesophageal problems, and 44.9% had renal disease (using the Cockcroft-Gault equation to calculate the glomerular filtration rate (GFR), with renal disease defined as an estimated GFR of less than 60 mL/min per 1.73m²).

CONCLUSIONS

Medication is an important part of osteoporosis treatment and fracture prevention but even in those whose doctor had told them they had osteoporosis, opportunities to use appropriate medications were missed and inappropriate medications were used.